

Opinion

Soil-Transmitted Helminthiasis and Vitamin A Deficiency: Two Problems, One Policy

Eric C. Strunz,¹ Parminder S. Suchdev,^{2,3} and David G. Addiss^{1,*}

Vitamin A deficiency (VAD) and soil-transmitted helminthiasis (STH) represent two widely prevalent and often overlapping global health problems. Approximately 75% of countries with moderate or severe VAD are coendemic for STH. We reviewed the literature on the complex relationship between STH and VAD. Treatment for STH significantly increases provitamin A (e.g., β -carotene) levels but is associated with minimal increases in preformed vitamin A (retinol). Interpretation of the data is complicated by variations in STH infection intensity and limitations of vitamin A biomarkers. Despite these challenges, increased coordination of STH and VAD interventions represents an important public health opportunity.

VAD and STH: Two Overlapping Global Health Problems

VAD and STH represent two widely prevalent and often overlapping global health problems. Affecting the world's poorest children and responsible for significant health and economic deficits, VAD and STH have been targeted by large-scale public health interventions for decades [1–3]. Recently, the desire to align campaigns for VAD and STH has grown, driven by the potential to efficiently share delivery platforms and harness synergistic health benefits from coadministering deworming drugs with vitamin A supplementation (VAS) [4,5]. While some studies have examined the impact of vitamin A status on STH [6–8], this review explores the effect of STH on vitamin A status.

Substantial geographic overlap exists between VAD and STH in preschool-age children. An estimated 75% (93/123) of countries with moderate or severe VAD also have preschool-age children who are considered by the World Health Organization (WHO) to be at risk of STH and in need of periodic mass treatment, known as 'preventive chemotherapy' [9]. Of the 190 million children affected by VAD globally, over 95% (~181 million) live in countries where the WHO recommends mass treatment for STH [9,10] (http://www.who.int/neglected_diseases/preventive_chemotherapy/sth/en/). Figure 1 (Key Figure) displays the overlap of countries with moderate or severe VAD and where preventive chemotherapy is required for STH.

The relationship between STH and vitamin A status remains unclear, however, as do the health advantages realized by coadministering treatments. Several studies have documented adverse effects of STH on vitamin A status but others have failed to detect a significant impact [11]. Interpreting the mixed body of evidence is further complicated by the challenges involved in measuring vitamin A status and STH.

This opinion article explores the nuanced relationship between STH and vitamin A. Relevant background on both conditions is first provided, followed by closer inspection of important

Trends

Available evidence suggests that soil-transmitted helminthiasis (STH) can contribute to vitamin A deficiency (VAD), although evidence on the impact of treatment for STH on VAD is inconclusive.

About 75% of countries with moderate or severe VAD are coendemic for STH.

Coordination of public health efforts to improve vitamin A status and treat STH provides synergies and public health benefits.

Expanding partnership between deworming and vitamin A supplementation efforts could help scale up deworming for preschool-age children.

¹Children Without Worms, The Task Force for Global Health, Decatur, GA, USA

²Department of Pediatrics, Emory University School of Medicine, Atlanta, GA, USA

³Hubert Department of Global Health, Rollins School of Public Health, Emory University, Atlanta, GA, USA

*Correspondence: daddiss@taskforce.org (D.G. Addiss).

Key Figure

Global Overlap of Vitamin A Deficiency and Soil-Transmitted Helminthiasis (STH)




Figure 1. Shaded countries are those that require preventive chemotherapy for STH according to the WHO. Cross-hatched countries are those with moderate or severe vitamin A deficiency (VAD) according to the WHO (i.e., prevalence of VAD is 10% or higher).

methodological considerations for assessment. Next we provide a review of the key literature before discussing implications for policymakers and public health practitioners. While additional research is needed to clarify the impact of STH on vitamin A status, coordination of campaigns for the two conditions could yield greater health benefits at lower costs than if implemented separately.

Vitamin A: An Essential Micronutrient

VAD, a condition resulting from inadequate vitamin A absorption, represents a significant contributor to mortality and morbidity for children and pregnant women, with a global burden estimated at 806 000 disability-adjusted life years (DALYs) [12]. Vitamin A is an essential micronutrient needed for vision, immune function, and cell differentiation [13]. Deficiency of this nutrient can cause xerophthalmia (i.e., dryness of the eyes), which can lead to night blindness and corneal ulceration [14].

VAD is commonly assessed by measuring the concentration of retinol in blood serum or through clinical eye examinations (Table 1) [9]. The WHO recommends using the prevalence of low serum retinol in preschool-age children to define the severity of VAD as a public health program, with $\geq 10\%$ and $\geq 20\%$ considered moderate and severe public health problems, respectively. Indicators of vitamin A have been recently reviewed by the Biomarkers of Nutrition for Development (BOND) program [15]. High-performance liquid chromatography (HPLC) is most frequently used to assess retinol concentrations; values $< 0.70 \mu\text{mol/l}$ indicate deficiency and $< 0.35 \mu\text{mol/l}$ represents severe deficiency. Nearly all of the circulating retinol in the blood is associated with its carrier protein, retinol-binding protein (RBP). Therefore, serum RBP can be used as a surrogate measure for retinol concentration [16]. Ocular examinations reveal a range of signs of VAD, including Bitot's spots, night blindness, and impaired dark adaptation [17]. Vitamin A indicators have been reviewed in the literature [17,18].

Table 1. Vitamin A Status Indicator Summary^a

Indicator	Assay	Classifications	Limitations
Serum retinol	HPLC	<0.70 $\mu\text{mol/l}$ is considered VAD <0.35 $\mu\text{mol/l}$ is considered severe VAD	Serum retinol levels are not a reliable indicator of the vitamin A liver stores Serum retinol levels show limited responsiveness to VAS
Dose response	RDR MRDR	RDR: 20% difference or greater in serum retinol concentrations pre- and post-serum retinol administration suggests low liver reserves of retinol MRDR: higher MRDR values suggest lower retinol reserves; a ratio of 3,4-didehydroretinol to retinol greater than 0.06 usually suggests low liver reserves of retinol	Cannot detect high (toxic) levels of vitamin A
Clinical signs	Visual eye inspection	Xerophthalmia can be evident from Bitot's spots (less severe) to blindness from corneal scarring (more severe); night blindness also indicates VAD	Ocular symptoms typically present only in cases of moderate-to-severe VAD
Liver sample	Liver biopsy	<0.07 $\mu\text{mol/g}$ liver of vitamin A represents deficient reserves	Gold standard for assessing vitamin A status Too invasive for public health studies
Isotope dilution	GCMS GCCIRMS	<0.07 $\mu\text{mol/g}$ liver of vitamin A represents deficient reserves	Optimal modern standard for assessing vitamin A status Not currently feasible for public health studies due to high time and cost requirements
Serum β -carotene	HPLC	No standard thresholds established	As a measure of provitamin A, β -carotene may be useful when assessing improvements to diet
Breast milk retinol	HPLC	$\leq 1.05 \mu\text{mol/l}$ (or $\leq 8 \text{ mg/g}$ milk fat) indicative of low breast-milk vitamin A content	Only applicable for lactating women Modest response to VAS

^aRDR, relative dose response; MRDR, modified relative dose response; GCMS, conventional gas chromatography–mass spectrometry; GCCIRMS, gas chromatography–combustion–isotope ratio mass spectrometry.

To help improve vitamin A status and reduce the burden of VAD, diet diversification and VAS have emerged as important public health interventions. VAS is typically administered orally using gelatin capsules containing 50 000–200 000 international units of vitamin A, most commonly to preschool-age children and women of childbearing age [19]. Diet diversification efforts have considered a range of strategies and foodstuffs, often adjusted for local context [20–23].

Although great progress has been achieved in recent decades, VAD remains an important cause of blindness and morbidity [24]. In 2009, the WHO estimated that night blindness affected over 5 million preschool-age children and almost 10 million pregnant women. Low serum retinol concentration (<0.70 $\mu\text{mol/l}$) was estimated to affect 190 million preschool-age children and 19 million pregnant women [9]. The geographic scope of VAD is also immense. Based on night blindness and serum retinol, 45 and 122 countries respectively suffer from VAD as a significant public health problem [9].

STH: A Disease of Poverty and the Environment

More than 1 billion people are at risk of STH, a group of diseases caused by the intestinal worms *Ascaris lumbricoides* (roundworm), *Trichuris trichiura* (whipworm), hookworm (*Ancylostoma duodenale* and *Necator americanus*), and *Strongyloides stercoralis* (threadworm) [4,25]. The

global burden of STH is estimated at over 5 million DALYs, largely due to anemia, stunting, and impaired cognitive development [12,26–28]. STH remains prevalent worldwide and represents a public health problem in 106 countries [10].

Humans are infected after ingesting eggs (*A. lumbricoides* and *T. trichiura*) or through penetration of the skin by larvae in the soil (hookworm) [28]. Current control strategies have focused on preventive chemotherapy, particularly for school-age children [29]. At-risk populations receive anthelmintic agents once or twice per year, usually a single oral dose of albendazole (400 mg) or mebendazole (500 mg). Treatments are typically provided in school settings but can also be delivered through community-wide initiatives or complementary distribution platforms (e.g., child-health days) [30]. Utilizing the ‘platform’ of VAS to coadminister deworming medicine could reduce costs and help scale up anthelmintic drug coverage among preschool-age children [31].

STH is typically diagnosed by microscopic examination of fecal samples for helminth eggs and the WHO recommends using the Kato–Katz method [29]. Other methods, including mini-FLOTAC and molecular diagnostics using PCR, have also been utilized, each with specific advantages. A recent review of diagnostics for STH provides an overview of the performance of the different assays [32]. Methodological considerations regarding the diagnosis of STH are summarized in Table 2.

Infections are categorized into three levels of intensity – light, moderate, and heavy – based on the number of eggs per gram of feces (epg). Higher egg counts, which correspond to more intense worm burdens, are associated with greater morbidity [29].

Mechanisms to explain the adverse effects of STH infection on nutrition have been proposed and include decreased appetite and nutrient intake, impaired absorption, and increased nutrient loss or altered metabolism [4]. Specifically for VAD, *Ascaris* expresses a range of retinol-binding proteins, retinol dehydrogenases, and retinoic acid receptors that utilize retinol for growth and

Table 2. Methodological Considerations Regarding the Diagnosis of STH

Consideration	Impact	Refs
Infection intensity	STH status is often reported as simply ‘infected’ or ‘not infected’, but the intensity of the worm burden represents a more meaningful indicator Morbidity is linked with intensity of infection and many individuals with light infections do not experience obvious symptoms Most individuals in an endemic area harbor light infections, while a small proportion are heavily infected and may harbor a large proportion of the overall community worm burden Many diagnostic methods, including the Kato–Katz method, will vary in sensitivity depending on intensity of infection and species of STH	[28,32–38]
Different helminth species	The term ‘soil-transmitted helminthiasis’ groups together infections from multiple, distinct intestinal nematodes that share the capability to be transmitted through soil <i>Ascaris lumbricoides</i> , <i>Trichuris trichiura</i> , <i>Ancylostoma duodenale</i> , <i>Necator americanus</i> , and <i>Strongyloides stercoralis</i> can differ in infection sequelae, susceptibility to current drug regimens, ease of detection, and other factors	[28,32,39,40]
Drug efficacy	Efficacy varies among different anthelmintic drugs; even a single drug will vary in efficacy against different helminths Albendazole, the most commonly used drug, yields fecal egg count reduction (FECR) rates of 99.5% for <i>A. lumbricoides</i> , 94.8% for hookworm, and 50.8% for <i>T. trichiura</i> Mebendazole yields FECR rates that also differ by helminth: 97.6% for <i>A. lumbricoides</i> , 79.6% for hookworm, and 63.1% for <i>T. trichiura</i> <i>S. stercoralis</i> infection can be particularly difficult to treat in a population context, as multiple doses or drug combinations (e.g., with ivermectin) are typically used to improve outcomes	[39–42]

development [43]. The pathways through which this occurs and their impacts on the human host remain unclear. Intestinal worms, due largely to their small biomass, consume only a small proportion of their host's dietary nourishment [44]. Other mechanisms may be more deleterious for nutrition, including helminth-induced tissue damage [45], decrease in host appetite [46], or simply increased metabolism and caloric demand resulting from the host's immune response [47]. Further, *A. lumbricoides* infection can alter the structure of the intestinal mucosa [48] and is associated with reduced fat absorption in humans [49,50] and animal models [51–54]. One trial randomized vitamin-A-deficient pigs to be infected by *Trichuris suis* and *Ascaris suum* (or serve as uninfected controls) and did not find a significant effect on either liver or serum retinol levels [55].

Inconclusive Evidence of STH Impact on Vitamin A Status

Evidence linking STH with vitamin A status is mixed. De Gier and colleagues recently conducted a meta-analysis that reviewed evidence linking helminth infections in school-age children with a range of micronutrient indicators [11]. Their comprehensive review included both experimental and observational studies of schistosomiasis or STH and numerous micronutrient markers (retinol, iron/ferritin, iodine, zinc, folate, and vitamin B12). Four randomized controlled trials (RCTs) were identified but none detected a significant difference in vitamin A status between treatment groups, which received anthelmintic agents, and control groups. A meta-analysis of 15 additional observational studies yielded a negative association between helminth infection and serum retinol (standardized mean difference: -0.30 ; 95% confidence interval: -0.48 , -0.13) [11]. However, the observational studies exhibited strong heterogeneity, suggesting that the meta-analysis averaged multiple distinct effects rather than describing the same particular relationship between helminth infection and serum retinol.

A recent study not included in the meta-analysis by de Gier and colleagues found a significant connection between STH and VAD. In periurban slums in Kenya, both any STH infection and *A. lumbricoides* infection were associated with a greater prevalence of VAD after controlling for demographic factors and nutritional status [56].

RCTs have yielded inconclusive findings regarding the intersection of vitamin A status and STH. Tanumihardjo and colleagues conducted a randomized trial exploring the relative effects of albendazole and VAS among enrolled Indonesian children ($n = 309$, 0.6–6.6 years old) with *A. lumbricoides* infection [57]. Sixty percent of enrolled children had light infections (1–4999 ep μ g), 30% had moderate infections (5000–49 999 ep μ g), and 10% had heavy infections ($\geq 50\,000$ ep μ g). Children were randomized into six treatment regimens, including a control group, to test the separate and combined effects of deworming and VAS. Across the experimental groups, deworming alone did not have a significant effect on the dehydroretinol/retinol ratio ($P = 0.37$) and the statistical interaction of treatment with albendazole and vitamin A was also not significant ($P = 0.75$).

Another randomized trial by Reddy *et al.* [58] also investigated the additive impact of deworming on VAS and found nonsignificant results. The authors investigated a population of 487 children (age 1–5 years) with 35% *A. lumbricoides* prevalence. They found no difference in serum retinol levels for children who received VAS compared with those who received deworming medications (tetramisole) and VAS. Deworming alone also did not improve vitamin A status compared with placebo. While serum retinol can be insensitive to changes in vitamin A status, both experimental groups that received VAS showed significantly increased serum retinol levels. This suggests that the null findings for deworming were not solely caused by the use of serum retinol as the outcome measure.

Another study by Mwaniki *et al.* [59] investigated the effects of multimicronutrient supplementation and multihelminth chemotherapy on serum retinol in Kenyan schoolchildren. In a four-arm

placebo-controlled randomized study, Mwaniki and colleagues investigated the effects of daily multimicronutrient supplementation over 8 months, with or without initial treatment using single-dose albendazole (for STH) and praziquantel (for schistosomiasis), in 977 Kenyan schoolchildren (age 7–19 years). Vitamin A supplementation increased serum retinol levels; praziquantel, but not albendazole, enhanced this effect. Prevalence was 14% for *A. lumbricoides*, 45% for *T. trichiura*, 54% for hookworm, and 71% for *Schistosoma mansoni*. Although egg counts were not reported, it is likely that all *Ascaris* infections were of low intensity due to the low prevalence.

Studies measuring preformed vitamin A (e.g., serum retinol) as the primary indicator of vitamin A status in relation to STH have found minimal significant results, but there is evidence to suggest that deworming can improve absorption of provitamin A, specifically β -carotene. Haque *et al.* conducted a four-arm, placebo-controlled randomized trial in Bangladesh investigating β -carotene supplementation and deworming in 244 preschool children who were infected with *A. lumbricoides* [60]. Children received low-dose β -carotene capsules daily for 6 months, either with or without albendazole, which was provided at the beginning of the study and 4 months after. Serum β -carotene increased by 0.08 $\mu\text{mol/l}$ after deworming, 0.06 $\mu\text{mol/l}$ after β -carotene supplementation, and 0.21 $\mu\text{mol/l}$ when treatments were combined. The increase in serum β -carotene measured in the group that received both deworming and supplementation was greater than the sum of the effects measured in both of the other groups, suggesting that a synergistic effect may be occurring with combined treatment.

There is also evidence that deworming may increase the effectiveness of food-based vitamin A interventions. A randomized study by Jalal *et al.* found that deworming improved serum retinol levels when combined with added dietary fat [61]. The combined effects of β -carotene-rich food, dietary fat, and deworming were also additive for participants with high-intensity infections. As a limitation, this study did not assess deworming separately from fat or food improvements, and those complementary interventions could confound the impact of deworming.

Beyond intervention studies, Ahmed and Mohiduzzaman did not find strong evidence linking ascariasis with malabsorption of vitamin A in humans [62]. In a study of 29 children in Bangladesh, they found that over 99% of vitamin A from supplements was absorbed successfully by the children. Only a small amount of retinol was recovered from stool samples. Expelled worms were also separated from stools and assayed, but no retinol was detected. However, there was a statistically significant negative linear relationship between the amount of vitamin A recovered from stools and worm burden. Most children in the study harbored light-intensity infections.

Concluding Remarks and Future Directions

Preformed vitamin A (i.e., retinol) has exhibited minimal or inconclusive responsiveness to deworming, but provitamin A (e.g., β -carotene) levels have increased significantly after treatment for STH. Considerations about the sensitivity and validity of serum retinol as an indicator for vitamin A status (Table 1) and the impact of STH infection intensity on disease morbidity make it difficult to interpret null findings from some trials. However, Tanumihardjo and colleagues largely account for these issues in their 1996 trial and provide the strongest evidence that treatment for STH may not improve vitamin A status directly [58]. Other motivations beyond improving vitamin A uptake may still justify coadministration of VAS with deworming.

In two studies that considered β -carotene as an indicator, Haque *et al.* and Jalal *et al.* found promising results after deworming [60,61]. As there is increased investment in sustainable dietary diversification efforts (e.g., promoting consumption of leafy green vegetables or orange-fleshed sweet potatoes) as well as mass-scale vitamin A food fortification (e.g., sugar, oil), the impact of alternative vitamin A interventions will need to be evaluated. Food fortification has

Outstanding Questions

What impact would integrated interventions have if implemented during early child development?

What is the impact of alternative vitamin A interventions, such as dietary diversification and vitamin A fortification, on vitamin A status in STH-endemic areas?

To what extent does coadministering treatment for STH and VAS increase coverage of VAS and save costs?

How should integrated STH and VAD programs be optimally monitored and evaluated?

proven to be a cost-effective strategy to address micronutrient deficiencies in developed countries and has been discussed as an important strategy for developing countries [63–66]. As evidenced by these two trials, dietary diversification to improve VAD could benefit from coadministered deworming.

Dosing of vitamin A could account for some of the discrepancy observed between the effects of STH on preformed vitamin A and provitamin A. Food fortification or diet diversification typically involve low, regular doses of vitamin A or provitamin A. High-dose supplementation with preformed vitamin A may interact differently with helminth infections. Further, most currently available data consider older, school-age children who may be less susceptible to morbidity from VAD compared with their younger peers. Additional research could examine the importance of early integrated interventions that target preschool-age children.

Other considerations may also support the value of adding deworming to current VAS efforts. No evidence has emerged indicating additional adverse effects caused by combining albendazole and vitamin A treatments. Even the large cluster-randomized Deworming and Enhanced Vitamin A (DEVTA) study found no evidence of immediate adverse events from albendazole and vitamin A treatment [67]. Further, deworming can improve participation in coadministered interventions since the results of deworming are readily observed in the stool [57]. Anthelmintic drugs appear to be popular in many communities, boosting treatment compliance and potentially increasing overall coverage of health interventions [4]. Community health volunteers also have reported that distributing deworming tablets strengthens their recognition in the community [5]. VAS cannot typically provide such clearly visible results, except in cases of xerophthalmia. Thus, deworming may strengthen the programmatic effectiveness of VAS without adding additional risks. Additional high-quality research would help facilitate a more thorough exploration of the synergy achieved by complementing VAS with deworming.

The WHO has established a target of regular deworming for at least 75% of at-risk preschool-age children by 2020 – an estimated 266 million children worldwide. In 2013, the most recent year for which data are available, global coverage as reported through the standard WHO channels was only 24% [10]. However, when UNICEF-assisted treatments coadministered with VAS were included, reported coverage increased to 49% [68]. This recent experience demonstrates the magnitude of coadministration already under way and the importance of the VAS platform for achieving the WHO deworming targets for preschool-age children.

While additional research is needed to clarify the relationship between soil-transmitted helminthiasis and vitamin A status, current evidence suggests that increasing coordination of campaigns for the two conditions represents an important opportunity that can yield significant public health benefits. However, based on existing evidence, program managers should not anticipate a significant increase in vitamin A uptake following the addition of deworming to VAS campaigns.

References

- Norris, J. *et al.* (2012) *Social and Economic Impact Review on Neglected Tropical Diseases. Economic Policy Briefing Paper*, Hudson Institute
- Sommer, A. (2008) Vitamin A deficiency and clinical disease: an historical overview. *J. Nutr.* 138, 1835–1839
- Bundy, D.A.P. *et al.* (2013) Worms, wisdom, and wealth: why deworming can make economic sense. *Trends Parasitol.* 29, 142–148
- Albonico, M. *et al.* (2008) Controlling soil-transmitted helminthiasis in pre-school-age children through preventive chemotherapy. *PLoS Negl. Trop. Dis.* 2, e126
- WHO (2004) *How to Add Deworming to Vitamin A Distribution*, WHO
- Long, K.Z. *et al.* (2007) Effect of vitamin A and zinc supplementation on gastrointestinal parasitic infections among Mexican children. *Pediatrics* 120, e846–e855
- Payne, L.G. *et al.* (2007) Benefit of vitamin A supplementation on *Ascaris* reinfection is less evident in stunted children. *J. Nutr.* 137, 1455–1459
- Al-Mekhlafi, H.M. *et al.* (2014) Does vitamin A supplementation protect schoolchildren from acquiring soil-transmitted helminthiasis? A randomized controlled trial. *Parasit. Vectors* 7, 367

9. WHO (2009) *Global Prevalence of Vitamin A Deficiency in Populations at Risk 1995–2005: WHO Global Database on Vitamin A Deficiency*, WHO
10. WHO (2015) Soil-transmitted helminthiasis: number of children treated in 2013. *Wkly. Epidemiol. Rec.* 90, 89–94
11. de Gier, B. *et al.* (2014) Helminth infections and micronutrients in school-age children: a systematic review and meta-analysis. *Am. J. Clin. Nutr.* 99, 1499–1509
12. Murray, C.J.L. *et al.* (2012) Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380, 2197–2223
13. Bailey, R.L. *et al.* (2015) The epidemiology of global micronutrient deficiencies. *Ann. Nutr. Metab.* 66 (Suppl. 2), 22–33
14. Black, R.E. (2014) Global distribution and disease burden related to micronutrient deficiencies. *Nestle Nutr. Inst. Workshop Ser.* 78, 21–28
15. Reiten, D.J. *et al.* (2011) Executive summary — biomarkers of nutrition for development: building a consensus. *Am. J. Clin. Nutr.* 94, 633S–650S
16. Erhardt, J.G. *et al.* (2004) Combined measurement of ferritin, soluble transferrin receptor, retinol binding protein, and C-reactive protein by an inexpensive, sensitive, and simple sandwich enzyme-linked immunosorbent assay technique. *J. Nutr.* 134, 3127–3132
17. Tanumihardjo, S.A. (2011) Vitamin A: biomarkers of nutrition for development. *Am. J. Clin. Nutr.* 94, 658S–665S
18. Rice, A.L. *et al.* (2000) Evaluation of serum retinol, the modified-relative-dose-response ratio, and breast-milk vitamin A as indicators of response to postpartum maternal vitamin A supplementation. *Am. J. Clin. Nutr.* 71, 799–806
19. WHO/UNICEF/IVACG Task Force *et al.* (1997) *Vitamin A Supplements: A Guide to Their Use in the Treatment and Prevention of Vitamin A Deficiency and Xerophthalmia*, WHO
20. Parlesak, A. *et al.* (2014) Toward the prevention of childhood undernutrition: diet diversity strategies using locally produced food can overcome gaps in nutrient supply. *Food Nutr. Bull.* 35, 191–199
21. Gurmu, F. *et al.* (2014) The potential of orange-fleshed sweet potato to prevent vitamin A deficiency in Africa. *Int. J. Vitam. Nutr. Res.* 84, 65–78
22. Hotz, C. *et al.* (2012) A large-scale intervention to introduce orange sweet potato in rural Mozambique increases vitamin A intakes among children and women. *Br. J. Nutr.* 108, 163–176
23. Bouis, H.E. *et al.* (2011) Biofortification: a new tool to reduce micronutrient malnutrition. *Food Nutr. Bull.* 32, 31S–40S
24. Sommer, A. (2014) Preventing blindness and saving lives: the centenary of vitamin A. *JAMA Ophthalmol.* 132, 115–117
25. Pullan, R.L. *et al.* (2014) Global numbers of infection and disease burden of soil transmitted helminth infections in 2010. *Parasit. Vectors* 7, 37
26. Chan, M-S. (1997) The global burden of intestinal nematode infections – fifty years on. *Parasitol. Today* 13, 438–443
27. Brooker, S. (2010) Estimating the global distribution and disease burden of intestinal nematode infections: adding up the numbers – a review. *Int. J. Parasitol.* 40, 1137–1144
28. Bethony, J. *et al.* (2006) Soil-transmitted helminth infections: ascariasis, trichuriasis, and hookworm. *Lancet* 367, 1521–1532
29. WHO (2011) *Helminth Control in School-Age Children: A Guide for Managers of Control Programmes*, WHO
30. Pan American Health Organization (2015) *Operational Guidelines for the Implementation of Deworming Activities: A Contribution to the Control of Soil-Transmitted Helminth Infections in Latin America and the Caribbean*, Pan American Health Organization
31. Rajagopal, S. *et al.* (2014) Micronutrient supplementation and deworming in children with geohelminth infections. *PLoS Negl. Trop. Dis.* 8, e2920
32. Nikolay, B. *et al.* (2014) Sensitivity of diagnostic tests for human soil-transmitted helminth infections: a meta-analysis in the absence of a true gold standard. *Int. J. Parasitol.* 44, 765–774
33. Larocque, R. *et al.* (2005) Relationship between intensity of soil-transmitted helminth infections and anemia during pregnancy. *Am. J. Trop. Med. Hyg.* 73, 783–789
34. Mascarini-Serra, L. (2011) Prevention of soil-transmitted helminth infection. *J. Glob. Infect. Dis.* 3, 175–182
35. Gabrielli, A.F. *et al.* (2011) Preventive chemotherapy in human helminthiasis: theoretical and operational aspects. *Trans. R. Soc. Trop. Med. Hyg.* 105, 683–693
36. Montresor, A. *et al.* (2015) *Trans. R. Soc. Trop. Med. Hyg.* 109, 262–267. <http://dx.doi.org/10.1093/trstmh/tru180>
37. Hall, A. *et al.* (1999) The distribution of *Ascaris lumbricoides* in human hosts: A study of 1765 people in Bangladesh. *Trans. R. Soc. Trop. Med. Hyg.* 93, 503–510
38. Bundy, D.A.P. (1988) Population ecology of intestinal helminth infections in human communities. *Phil. Trans. R. Soc. Lond. B* 321, 405–442
39. Levecke, B. *et al.* (2014) Assessment of anthelmintic efficacy of mebendazole in school children in six countries where soil-transmitted helminths are endemic. *PLoS Negl. Trop. Dis.* 8, e3204
40. Vercruysse, J. *et al.* (2011) Assessment of the anthelmintic efficacy of albendazole in school children in seven countries where soil-transmitted helminths are endemic. *PLoS Negl. Trop. Dis.* 5, e948
41. Pungpak, S. *et al.* (1987) Albendazole in the treatment of strongyloidiasis. *Southeast Asian J. Trop. Med. Public Health* 18, 207–210
42. Archibald, L.K. *et al.* (1993) Albendazole is effective treatment for chronic strongyloidiasis. *Q. J. Med.* 86, 191–195
43. Hurst, R.J.M. and Else, K.J. (2012) Retinoic acid signalling in gastrointestinal parasite infections: lessons from mouse models. *Parasite Immunol.* 34, 351–359
44. Hall, A. *et al.* (2008) A review and meta-analysis of the impact of intestinal worms on child growth and nutrition. *Matern. Child. Nutr.* 4, 118–236
45. Hotez, P.J. and Cerami, A. (1983) Secretion of a proteolytic anticoagulant by *Ancylostoma hookworms*. *J. Exp. Med.* 157, 1594–1603
46. Hadju, V. *et al.* (1996) Improvements in appetite and growth in helminth-infected schoolboys three and seven weeks after a single dose of pyrantel pamoate. *Parasitology* 113, 497–504
47. Hall, A. (1985) Nutritional aspects of parasitic infection. *Prog. Food Nutr. Sci.* 9, 227–256
48. Crompton, D.W.T. and Nesheim, M.C. (2002) Nutritional impact of intestinal helminthiasis during the human life cycle. *Annu. Rev. Nutr.* 22, 35–59
49. Sivakumar, B. and Reddy, V. (1975) Absorption of vitamin A in children with ascariasis. *J. Trop. Med. Hyg.* 78, 114–115
50. Mahalanabis, D. *et al.* (1976) Vitamin A absorption in ascariasis. *Am. J. Clin. Nutr.* 29, 1372–1375
51. Stephenson, L.S. *et al.* (1980) *Ascaris suum*: nutrient absorption, growth, and intestinal pathology in young pigs experimentally infected with 15-day-old larvae. *Exp. Parasitol.* 49, 15–25
52. Forsum, E. *et al.* (1981) Nutritional aspects of *Ascaris* infection in young protein-deficient pigs. *Parasitology* 83, 497–512
53. Tripathy, K. *et al.* (1971) Effects of *Ascaris* infection on human nutrition. *Am. J. Trop. Med. Hyg.* 20, 212–218
54. Brown, K.H. *et al.* (1980) Absorption of macronutrients from a rice-vegetable diet before and after treatment of ascariasis in children. *Am. J. Clin. Nutr.* 33, 1975–1982
55. Pedersen, S. *et al.* (2001) Marginal vitamin A deficiency in pigs experimentally infected with *Trichuris suis*: a model for vitamin A inadequacy in children. *Trans. R. Soc. Trop. Med. Hyg.* 95, 557–565
56. Suchdev, P.S. *et al.* (2014) Soil-transmitted helminth infection and nutritional status among urban slum children in Kenya. *Am. J. Trop. Med. Hyg.* 90, 299–305
57. Tanumihardjo, S.A. *et al.* (1996) Vitamin A status of Indonesian children infected with *Ascaris lumbricoides* after dosing with vitamin A supplements and albendazole. *J. Nutr.* 126, 451–457
58. Reddy, V. *et al.* (1986) Effect of deworming and vitamin A administration on serum vitamin A levels in preschool children. *J. Trop. Pediatr.* 32, 196–199
59. Mwaniki, D. *et al.* (2002) Effects on serum retinol of multi-micronutrient supplementation and multi-helminth chemotherapy: a

- randomised, controlled trial in Kenyan school children. *Eur. J. Clin. Nutr.* 56, 666–673
60. Haque, R. *et al.* (2010) Low-dose β -carotene supplementation and deworming improve serum vitamin A and β -carotene concentrations in preschool children of Bangladesh. *J. Health Popul. Nutr.* 28, 230–237
 61. Jalal, F. *et al.* (1998) Serum retinol concentrations in children are affected by food sources of β -carotene, fat intake, and anthelmintic drug treatment. *Am. J. Clin. Nutr.* 68, 623–629
 62. Ahmed, F. *et al.* (1993) Vitamin A absorption in children with ascariasis. *Br. J. Nutr.* 69, 817–825
 63. Fulgoni, V.L. and Buckley, R.B. (2015) The contribution of fortified ready-to-eat cereal to vitamin and mineral intake in the U.S. population, NHANES 2007–2010. *Nutrients* 7, 3949–3958
 64. Dary, O. and Mora, J.O. (2002) Food fortification to reduce vitamin A deficiency: International Vitamin A Consultative Group recommendations. *J. Nutr.* 132, 2927S–2933S
 65. Haas, J.H. and Miller, D.D. (2006) Overview of Experimental Biology 2005 Symposium: Food Fortification in Developing Countries. *J. Nutr.* 136, 1053–1054
 66. Fiedler, J.L. *et al.* (2015) Assessing alternative industrial fortification portfolios: a Bangladesh case study. *Food Nutr. Bull.* 36, 57–74
 67. Awasthi, S. *et al.* (2013) Population deworming every 6 months with albendazole in 1 million pre-school children in north India: DEVTA, a cluster-randomised trial. *Lancet* 381, 1478–1486
 68. Kumapley, R.S. *et al.* (2015) The role of child health days in the attainment of global deworming coverage targets among pre-school-age children. *PLoS Negl. Trop. Dis.* 9, e0004206